



Two roles for ecological surrogacy: Indicator surrogates and management surrogates



Malcolm Hunter Jr.^{a,*}, Martin Westgate^b, Philip Barton^b, Aram Calhoun^a, Jennifer Pierson^b, Ayesha Tulloch^b, Maria Beger^c, Cristina Branquinho^d, Tim Caro^e, John Gross^f, Jani Heino^g, Peter Lane^b, Catherine Longo^h, Kathy Martinⁱ, William H. McDowell^j, Camille Mellin^k, Hanna Salo^l, David Lindenmayer^b

^a Department of Wildlife, Fisheries, and Conservation Biology, University of Maine, Orono, ME 04649, USA

^b Fenner School of Environment and Society, The Australian National University, Acton, Australian Capital Territory 2601, Australia

^c Centre for Biodiversity and Conservation Science, The University of Queensland, Brisbane, Queensland 4072, Australia

^d Centre for Ecology, Evolution and Environmental Changes, Faculty of Sciences, University of Lisbon, 1749-016 Lisboa, Portugal

^e Department of Wildlife, Fish and Conservation Biology, University of California, Davis, CA 95616, USA

^f Climate Change Response Program, U.S. National Park Service, Fort Collins, CO 80525, USA

^g Finnish Environment Institute, Natural Environment Centre, Biodiversity, FI-90014 Oulu, Finland

^h National Center for Ecological Analysis and Synthesis, Santa Barbara, CA 93101, USA

ⁱ Department of Forest and Conservation Sciences, University of British Columbia, Vancouver, BC V6 T 1Z4, Canada

^j Department of Natural Resources and the Environment, University of New Hampshire, Durham, NH 03824, USA

^k Australian Institute of Marine Science, Townsville, Queensland 4810, Australia

^l Department of Geography and Geology, University of Turku, Turku, Finland

ARTICLE INFO

Article history:

Received 21 July 2015

Received in revised form

16 November 2015

Accepted 22 November 2015

Available online 23 December 2015

Keywords:

Coarse-filter

Environmental management

Flagship species

Focal species

Indicators

Monitoring

Environmental proxy

Surrogates

Terminology

Umbrella species

ABSTRACT

Ecological surrogacy – here defined as using a process or element (e.g., species, ecosystem, or abiotic factor) to represent another aspect of an ecological system – is a widely used concept, but many applications of the surrogate concept have been controversial. We argue that some of this controversy reflects differences among users with different goals, a distinction that can be crystallized by recognizing two basic types of surrogate. First, many ecologists and natural resource managers measure “indicator surrogates” to provide information about ecological systems. Second, and often overlooked, are “management surrogates” (e.g., umbrella species) that are primarily used to facilitate achieving management goals, especially broad goals such as “maintain biodiversity” or “increase ecosystem resilience.” We propose that distinguishing these two overarching roles for surrogacy may facilitate better communication about project goals. This is critical when evaluating the usefulness of different surrogates, especially where a potential surrogate might be useful in one role but not another. Our classification for ecological surrogacy applies to species, ecosystems, ecological processes, abiotic factors, and genetics, and thus can provide coherence across a broad range of uses.

© 2015 Elsevier Ltd. All rights reserved.

1. Introduction

In October 2014 a diverse group of scientists from around the world gathered in Australia to spend three days exploring the full scope of ecological surrogacy, primarily trying to achieve a broad, synthetic understanding that would advance the use of this

important concept. They ranged from conservation practitioners and scientists who use bacteria and lichens to monitor pollution, to those who try to foster ecological integrity of whole oceans, or try to conserve regional biodiversity by managing representative arrays of ecosystems. The participants soon discovered that, despite a common interest in the use of surrogates for monitoring and managing ecological systems, they did not share a foundational understanding of ecological surrogacy. In particular, those who measure surrogates as ecological indicators found it difficult to embrace the concept of surrogates as alternative foci for

* Corresponding author. Tel.: +1 2075812865.

E-mail address: mhunter@maine.edu (M. Hunter Jr.).

Table 1
Examples of indicator surrogates and management surrogates for five types of ecological components. The example goals highlight distinctions between monitoring and managing. We chose these five classes to show the wide applicability of our concept, not to suggest that they constitute a definitive classification (e.g., one could readily combine ecosystems and ecosystem processes or separate abiotic factors into chemical and physical factors).

Class	Indicator surrogates		Management surrogates	
	Example	Example goal	Example	Example goal
Species	Indicator species	Detect change in target species abundance (<i>E. coli</i> , Ashbolt et al., 2001)	Umbrella species	Conserve a large suite of species (<i>Leadbeater's possum</i> , Lindenmayer, 1996)
	Indicator guilds	Detect change in function provided by a guild (pollinators, Kehinde and Samways, 2012)	Flagship species	Foster support for conservation (giant panda, Bowen-Jones and Entwistle, 2002)
Ecosystems	Ecosystem extent	Use species–area relationships to predict species richness (Triantis et al., 2015)	Ecosystems as coarse filters	Maintain biodiversity at species and genetic level by conserving a representative array of ecosystems (Hunter et al., 1988)
	Ecosystem structure	Measure structural diversity to quantify habitat for target species (Baril et al., 2011)		
Ecological processes	Ecosystem productivity	Detect changes in biomass accumulation (Culman et al., 2010)	Disturbance regimes	Manage fire regimes to create desired vegetation (Bradstock et al., 2012)
	Biogeochemical cycling	Detect carbon fluxes (Fan et al., 2015)	River flows	Manage flow regimes to restore riverine ecosystem integrity (Beechie et al., 2010)
Abiotic factors	Nutrient concentration	Monitor nitrogen and phosphorous water pollution (Rocha et al., 2015)	Geological and climatic diversity	Conserve diverse environments for biodiversity (Beier et al., 2015)
Genetics	Population structure	Detect functional connectivity (Braunisch et al., 2010)	Genetic diversity	Maintain evolutionary potential (tuatara, Miller et al., 2012)

management. For example, managing an umbrella species because it is an efficient way to maintain a large set of species did not seem like a form of surrogacy to them, whereas this was a key form of surrogacy for others. This led to many hours of discussion and ultimately we reached a consensus that explicitly recognizes two basic forms of surrogates based on goals: indicator surrogates (which are measured to provide information about ecological systems) and management surrogates (which are managed to achieve a different, often larger, goal such as “maintain biodiversity”). In this paper, we argue that this dichotomy represents a meaningful division in how different groups use ecological surrogacy. We discuss why this schism has emerged, and give examples of how it applies to five types of ecological components: species, ecosystems, ecological processes, abiotic factors, and genetics. We argue that disagreements over surrogate utility typically occur between groups with different goals, and that by explicitly recognizing two overarching goals for ecological surrogacy – providing information about ecological systems and facilitating their management – future misunderstandings can be avoided.

2. Context and definitions

Although ecological surrogacy is a frequently used concept (nearly 50,000 journal articles by one count; [Westgate et al., 2014](#)), it has repeatedly defied simple classification. For example, the United Nations ([UNCCD 2013](#)), [Secretariat of the Convention on Biological Diversity \(2014\)](#), European Union ([BISE, 2014](#)), Australia’s State of the Environment Program ([ANZECC State of the Environment Reporting Task Force 2000](#)), and the United States Fish and Wildlife Service ([USFWS, 2014](#)) have all embraced different uses of surrogates. This lack of consensus amongst academics and practitioners on a shared terminology or scheme of classification remains despite repeated critiques and attempts at standardization (e.g., [Landres et al., 1988](#); [Noss, 1990](#); [McGeogh, 1998](#); [Simberloff, 1998](#); [Dale and Beyeler, 2001](#); [Niemi and McDonald, 2004](#); [Caro,](#)

[2010](#); [Heink and Kowarik, 2010](#); [Pereira et al., 2013](#); [Lindenmayer et al., 2015](#); [Stephens et al., 2015](#)). We propose to build a simple foundation for ecological surrogacy by recognizing that many seemingly distinct applications of the surrogate concept share common goals: environmental monitoring or informing management. Our focus on goals differs from earlier classification schemes that emphasized differences among organizational scales (e.g., genes, species, or ecosystems; [Caro, 2010, Table 1](#)), ecological attributes (e.g., compositional, functional, or structural; [Noss, 1990](#)), or distinct types of problem (e.g., environment, ecology, or biodiversity surrogates; [McGeogh, 1998](#)).

We begin with a definition of ecological surrogacy to distinguish it from surrogacy in medicine, engineering, and other fields ([Forrester et al., 2008](#); [Barton et al., 2014](#)).

Ecological surrogate: An ecological process or element (e.g., species, ecosystem, or abiotic factor) that is used to represent (i.e., serve as a proxy for) another aspect of an ecological system.

The earliest explicit uses of surrogates focused on measuring one species as an indicator for others: i.e., beginning in 1893, the concentration of *Escherichia coli* was used to indicate the likely presence of other pathogens in drinking water ([Ashbolt et al., 2001](#)). This usage is clearly consistent with our definition of indicator surrogates:

Indicator surrogate: A type of surrogate that provides information about another aspect of an ecological system: for example, measuring the population density of species A because it provides information about the condition of target ecosystem X.

This approach emphasizes a mechanistic, statistical approach to surrogacy that remains popular amongst environmental scientists. However, a dramatic expansion in the use of the surrogacy concept in ecology and conservation biology arose alongside the development of the concept of “biodiversity” in the 1980s. Advocates of maintaining biodiversity realized that it was impractical to address

directly all of the elements of biodiversity given the vast numbers of species, especially little-known or undescribed invertebrates and microbes, or the genetic components of biodiversity. Thus, conservation practitioners needed surrogates that could be readily managed under the assumption that managing the surrogate would be beneficial for a sizable portion of biodiversity. (In this paper we use “management” broadly to cover activities as diverse as controlling a contaminant, conserving a game species or endangered species, planning a reserve system, or motivating public support for conservation.) From this emerged the idea of a conservation “umbrella” in which one species is used to represent biodiversity for management purposes (Frankel and Soule, 1981). Similarly, but on an ecosystem level, “coarse-filter” conservation assumed that protecting a representative array of ecosystems would encompass much biodiversity at the species and even genetic levels, with relatively few species falling through the filter’s pores unprotected (Noss, 1987). With both umbrella species and coarse filters, the primary goal is to manage X to achieve the real target goal Y. In addition to biodiversity, other broad conceptual entities such as ecological integrity (Rapport et al., 1998) or resilience (Walker and Salt, 2012) have also become the basis for setting large goals that are often addressed using proxies that we call “management surrogates.”

Management surrogate: A type of surrogate that is a tool for management because it represents another aspect of an ecological system that is the main goal of management: for example, managing the population of species A because this facilitates maintaining the integrity of ecosystem X.

Therefore, management surrogacy focuses primarily on facilitating management of ecological systems whereas indicator surrogacy focuses primarily on providing information about those systems.

We suggest that our binary, goal-oriented approach to surrogate classification represents an improvement over existing schemes for two reasons. First, our conceptual understanding of how to classify ecological systems is diverse and evolving, but goals of measuring and managing ecosystems are relatively constant. Second, scientists often do not articulate clear, explicit goals, and so discussion about goal setting is likely to be beneficial to the science and application of surrogates.

The definitions provided above are distinct, but when surrogates are applied in practice there can easily be overlap; we turn to this issue next.

3. Divergent goals and surrogate effectiveness

When management surrogates and indicator surrogates are seen as complementary constructs, some past debates over surrogate effectiveness can be reinterpreted as differences between users with different goals and approaches (Caro, 2010). For example, controversies over the utility of focal species (Lambeck, 1997; Lindenmayer et al., 2002) or flagship species (Simberloff, 1998) may have arisen because management surrogates were incorrectly interpreted as indicator surrogates.

To illuminate the distinctions and overlaps between management and indicator surrogates, we offer three well-known examples. First, we consider an example in which a species might be an effective management surrogate for biodiversity even though it would probably be an ineffective indicator surrogate of biodiversity. Tigers (*Panthera tigris*) are difficult to count and select habitat at coarser scales than most species, and thus are an ineffective indicator surrogate. Yet organizing biodiversity management around tigers as an umbrella species may be sensible because conserving their habitat (mangrove swamps to boreal forests) would provide habitat for thousands of other species (Wikramanayake

et al., 2008). A converse example (an effective indicator surrogate that is an insufficient management surrogate) may be found with *E. coli*. Monitoring *E. coli* may indicate if rivers are free of fecal contamination, but in many rivers, the key to restoring ecological integrity is fostering natural river flows (e.g., removing in-stream barriers and managing flow through dams, Beechie et al., 2010). In this case, an ecological process, the flow regime, would be a more effective management surrogate. These two examples are clearcut, but when surrogates are applied in practice there is often substantial overlap between management and indicator surrogacy. Consider the role of beavers (*Castor canadensis* and *C. fiber*) in providing habitat for pond-dependent biota. If we monitor beaver populations with the goal of assessing and tracking habitat availability for other species such as waterfowl, beavers are serving as an *indicator surrogate*. If we increase the beaver population, perhaps by banning trapping, with the goal of increasing the number of beaver dams and thus ponds, this is *management surrogacy*. In many cases, these approaches will be coordinated and thus both forms of surrogacy used, but this is not necessarily the case. One could manage beavers to increase the number of beaver ponds without systematically monitoring their populations. Alternatively, one could monitor beavers to indicate changes in other pond-dwelling species, without any active beaver management.

4. Surrogacy in five classes of ecological components

To demonstrate how our definitions relate to surrogate use in practice, we apply the “indicator surrogates” and “management surrogates” concept to five classes of ecological components: species, ecosystems, ecological processes, abiotic factors, and genetics. We selected these to show the wide applicability of our concept, not to imply that they are the basis of a robust classification (e.g., one could readily combine ecosystem processes and ecosystems or split abiotic factors into physical and chemical factors). We begin with species because here the language is most developed (Table 1).

1. Surrogate species terminology is quite easy to distill into indicator surrogates and management surrogates because we have a well-established term, “indicator species”, clearly linked to measuring one component of an ecosystem to represent another component, as well as two common terms, umbrella species and flagship species, which are primarily linked to management (Caro, 2010). The indicator species concept has many different refinements (e.g., sentinel species, biomonitoring species, ecological-disturbance indicator species; Caro, 2010) and has been extended to include indicator taxa (e.g., lichens; Brunialti et al., 2009) and using species traits (Moretti and Legg, 2009). Simply counting species to estimate species richness is a commonly used indicator surrogate, sometimes employed to estimate the species richness of a different taxon, sometimes used to assess the status of an ecosystem (Fleishman et al., 2006). The management surrogate concept can also be extended from individual species to umbrella taxa (see the Important Bird Areas program; BirdLife International, 2004) and umbrella guilds (Drever et al., 2008), and it is related to other approaches for identifying species that might be particularly important for management, such as keystone species or landscape species (Caro, 2010).
2. Ecosystems, like species, have well established roles as both indicator surrogates and management surrogates, but the terminology is not as explicit. One rarely hears of indicator ecosystems, umbrella ecosystems, or flagship ecosystems, even though it could be argued that coral reefs and rainforests are flagships due to their public prominence. The areal extent of an ecosystem is

the most commonly used index of its indicator surrogacy value, although spatial configuration or connectivity are sometimes evaluated too. It is also common to measure ecosystem components such as vegetation structure (Noss, 1990) as indicators of overall condition. Ecosystems also have a critical role as management surrogates in the context of maintaining biodiversity at the species and genetic levels, i.e., the so-called coarse-filter strategy (Hunter et al., 1988), and thus conservation planning to maintain a representative array of ecosystems is well-established (Groves, 2003).

3. Ecological processes are commonly used for indicator surrogacy; in particular measurements of such key features as ecosystem productivity and biogeochemical cycling are used as indicator surrogates for ecosystem condition (Noss, 1990). Additionally, in recent decades, some processes that can be manipulated or even emulated have become management surrogates. For example, fire can be a management surrogate because ecosystem managers in fire-prone ecosystems often seek to maintain fire regimes that meet ecological and societal goals, including the provision of habitat for fire-dependent species (Bradstock et al., 2012). In forestry, the idea of emulating natural disturbance and succession regimes through specific timber management practices (Lindenmayer, 2009) has meant that these processes are used as management surrogates tied to larger goals such as biodiversity and ecological integrity (Hunter and Schmiegelow, 2011).
4. Abiotic factors are widely used as indicator surrogates; e.g., monitoring dissolved oxygen or pH to understand lake condition. It is also common for abiotic factors to be used as management surrogates. For example, when climate mitigation strategies are organized around reducing atmospheric CO₂, then CO₂ is a management surrogate for the much larger, more complex climate system. Management surrogacy centered on abiotic factors is the foundation of proposals to adapt to climate change by designing reserve systems around enduring abiotic factors such as topography, geology, and hydrology (Beier and Brost, 2010; Beier et al., 2015).
5. Genetic metrics have a steadily growing role as indicator surrogates through genetic monitoring (Schwartz et al., 2007), especially in relation to genetic erosion (Hoban et al., 2014), effective population size (Tallmon et al., 2012), and landscape connectivity (Baguette et al., 2013). Genetics also has a role as a management surrogate, particularly because maintaining genetic diversity is a means to achieve the larger goal of safeguarding evolutionary potential (Harrisson et al., 2014) and resilience (Schindler et al., 2010).

In summary, we recognize that both indicator surrogates and management surrogates are widely used across diverse components of ecological systems (Table 1). The use of indicator surrogates is more established, but the use of management surrogates is increasing in response to broad goals like maintaining ecosystem integrity and biodiversity.

5. Benefits of setting clearer goals

The main advantage of our proposed construct is that it gives explicit recognition of two overarching goals for ecological surrogacy that are linked but conceptually separate: to provide information about ecological systems and to facilitate their management. By seeing how this concept and related terms fit together (Table 1), it should be clearer how different disciplines might better learn from each other, and open up new opportunities for synthetic thinking and analysis. People as diverse as those who monitor lichen uptake of air pollutants (Brunialti et al., 2009) and

those who try to assess the ecosystem services of oceans (Halpern et al., 2012) need to speak a common language, or at least agree on some fundamental ideas to foster cross-disciplinary learning. It is particularly important to recognize that an imperfect indicator surrogate might still serve as a useful management surrogate, and vice versa. Furthermore, the science underpinning each kind of surrogate may not be transferable, and research on each should be framed and assessed in relation to specific explicit goals.

We have argued that disagreements over the utility of ecological surrogates may reflect a misalignment of the goals of people who use indicator surrogates versus management surrogates (see Westgate et al., 2013). Such differences might also reveal a schism in opinions about the value of quantitative information for improving conservation outcomes. As scientists, we have an implicit bias toward evidence-based approaches (e.g., Sutherland et al., 2004) and this may blind us to policy and public communication benefits of management surrogates that can be difficult to quantify. For example, promoting flagship or umbrella species can lead to large conservation gains, particularly if stakeholders are more likely to embrace a single charismatic species than a set of ecological metrics serving as indicator surrogates (Schultz, 2011). In short, disagreements over the utility of surrogates may reflect deeper arguments about the role of scientific information in conservation practice (Mace, 2014).

In conclusion, ecological surrogacy is widely used by natural resource management organizations around the world, and that usage will probably increase because of its potential expediency and efficiency. To avoid unproductive and circular debates, we have sought clarity by explicitly recognizing two different (but equally legitimate) core uses for ecological surrogacy. We argue that evidence of surrogate efficacy may be based on the success of a management program (e.g., increased public support following a flagship species campaign), or documentation of a tight ecological relationship between a surrogate and its target (e.g., linking population viability of a species to ecosystem integrity), or both. Recognizing that different stakeholders have different goals when using surrogates should foster communication and collaboration across a wide range of disciplines, and thus build a multi-disciplinary foundation for effective ecological management.

Acknowledgments

The work in this paper was funded by the Australian Research Council through a Laureate Fellowship to DBL. Tabitha Boyer and Claire Shepherd assisted with many aspects of workshop organization that brought together the co-authors of this article and subsequent manuscript preparation. Paul Beier, Bonnie Lamb, Lee O'Brien, and Greg Shriver reviewed earlier drafts of this manuscript.

References

- ANZECC (Australian and New Zealand Environment and Conservation Council) State of the Environment Reporting Task Force, 2000. Core Environmental Indicators for Reporting on the State of the Environment. Environment Australia, Canberra, Available from <http://www.environment.gov.au/node/23085> (accessed 03.08.14).
- Ashbolt, N.J., Grabow, W.O.K., Snozzi, M., 2001. Indicators of microbial water quality. In: Fewtrell, L., Bartram, J. (Eds.), *Water Quality: Guidelines, Standards and Health*. IWA Publishing, London, UK, pp. 289–316.
- Baguette, M., Blanchet, S., Legrand, D., Stevens, V.M., Turlure, C., 2013. Individual dispersal, landscape connectivity and ecological networks. *Biol. Rev.* 88, 310–326.
- Baril, L.M., Hansen, A.J., Renkin, R., Lawrence, R., 2011. Songbird response to increased willow (*Salix* spp.) growth in Yellowstone's northern range. *Ecol. Appl.* 21, 2283–2296.
- Barton, P.S., Pierson, J.C., Westgate, M.J., Lane, P.W., Lindenmayer, D.B., 2014. Learning from clinical medicine to improve the use of surrogates in ecology. *Oikos* 124, 391–398.
- Beechie, T.J., Sear, D.A., Olden, J.D., Pess, G.R., Buffington, J.M., Moir, H., Roni, P., Pollock, M.M., 2010. Process-based principles for restoring river ecosystems. *Bioscience* 60, 209–222.

- Beier, P., Brost, B., 2010. Use of land facets to plan for climate change: conserving the arenas, not the actors. *Conserv. Biol.* 24, 701–710.
- Beier, P., Hunter, M.L., Anderson, M., 2015. Conserving nature's stage. *Conserv. Biol.* 29, 613–617.
- BirdLife International, 2004. *State of the World's Birds 2004: Indicators for Our Changing World*. BirdLife International, Cambridge, UK.
- BISE, 2014. EU Biodiversity Indicators and Related Global Aichi Targets. Biodiversity Information System for Europe, Available from <http://biodiversity.europa.eu/policy/eu-biodiversity-indicators-and-related-global-aichi-targets> (accessed 23.11.14).
- Bowen-Jones, E., Entwistle, A., 2002. Identifying appropriate flagship species: the importance of culture and local contexts. *Oryx* 36, 189–195.
- Bradstock, R.A., Gill, M., Williams, R.J., 2012. *Flammable Australia: Fire Regimes, Biodiversity and Ecosystems in a Changing World*. CSIRO Publishing, Melbourne.
- Braunisch, V., Segelbacher, G., Hirzel, A.H., 2010. Modelling functional landscape connectivity from genetic population structure: a new spatially explicit approach. *Mol. Ecol.* 19, 3664–3678.
- Brunialti, G., Frati, L., Incerti, G., Rizzi, G., Vinci, M., Giordani, P., 2009. Lichen biomonitoring of air pollution: issues for applications in complex environments. In: Romano, G.C., Conti, A.C. (Eds.), *Air Quality in the 21st Century*. NOVA Science Publisher, Hauppauge, NY, pp. 1–48.
- Caro, T., 2010. *Conservation by Proxy: Indicator, Umbrella, Keystone, Flagship, and Other Surrogate Species*. Island Press, Washington, DC.
- Culman, S.W., Young-Mathews, A., Hollander, A.D., Ferris, H., Sánchez-Moreno, S., O'Geen, A.T., Jackson, L.E., 2010. Biodiversity is associated with indicators of soil ecosystem functions over a landscape gradient of agricultural intensification. *Landsc. Ecol.* 25, 1333–1348.
- Dale, V.H., Beyeler, S.C., 2001. Challenges in the development and use of ecological indicators. *Ecol. Indic.* 1, 3–10.
- Drever, M., Aitken, K., Norris, A., Martin, K., 2008. Woodpeckers as reliable indicators of bird richness, forest health and harvest. *Biol. Conserv.* 141, 624–634.
- Fan, L.-C., Yang, M.-Z., Han, W.-Y., 2015. Soil respiration under different land uses in Eastern China. *PLoS One* 10, e0124198.
- Fleishman, E., Noss, R.F., Noon, B.R., 2006. Utility and limitations of species richness metrics for conservation planning. *Ecol. Indic.* 6, 543–553.
- Forrester, A., Sobester, A., Keane, A., 2008. *Engineering Design via Surrogate Modelling: A Practical Guide*. John Wiley and Sons, Chichester, UK.
- Frankel, O.H., Soule, M.E., 1981. *Conservation and Evolution*. Cambridge University Press, Cambridge, UK.
- Groves, C., 2003. *Drafting a Conservation Blueprint*. Island Press, Washington, DC.
- Halpern, B.S., et al., 2012. An index to assess the health and benefits of the global ocean. *Nature* 488, 615–620.
- Harrisson, K.A., Pavlova, A., Telonis-Scott, M., Sunnucks, P., 2014. Using genomics to characterize evolutionary potential for conservation of wild populations. *Evol. Appl.* 7, 1008–1025.
- Heink, U., Kowarik, I., 2010. On the definition of indicators in ecology and environmental planning. *Ecol. Indic.* 10, 584–593.
- Hoban, S., Arntzen, J.A., Bruford, M.W., Godoy, J.A., Rus Hoelzel, A., Segelbacher, G., Vilà, C., Bertorelle, G., 2014. Comparative evaluation of potential indicators and temporal sampling protocols for monitoring genetic erosion. *Evol. Appl.* 7, 984–998.
- Hunter Jr., M.L., Schmiegelow, F., 2011. *Wildlife, Forests, and Forestry: Principles of Managing Forests for Biological Diversity*, 2nd ed. Prentice-Hall, Upper Saddle River, NJ.
- Hunter Jr., M.L., Jacobson Jr., G.L., Webb III, T., 1988. Paleoeecology and the coarse-filter approach to maintaining biological diversity. *Conserv. Biol.* 2, 375–385.
- Kehinde, T., Samways, M.J., 2012. Endemic pollinator response to organic vs. conventional farming and landscape context in the Cape Floristic Region biodiversity hotspot. *Agric. Ecosyst. Environ.* 146, 162–167.
- Lambeck, R.J., 1997. Focal species: a multi-species umbrella for nature conservation. *Conserv. Biol.* 11, 849–856.
- Landres, P.B., Verner, J., Thomas, J.W., 1988. Ecological uses of vertebrate indicator species: a critique. *Conserv. Biol.* 2, 316–328.
- Lindenmayer, D.B., 1996. *Wildlife and Woodchips: Leadbeater's Possum a Test Case of Sustainable Forestry*. UNSW Press, Sydney.
- Lindenmayer, D.B., 2009. *Forest Pattern and Ecological Process: A Synthesis of 25 Years of Research*. CSIRO Publishing, Melbourne.
- Lindenmayer, D.B., Manning, A.D., Smith, P.L., Possingham, H.P., Fischer, J., Oliver, I., McCarthy, M.A., 2002. The focal-species approach and landscape restoration: a critique. *Conserv. Biol.* 16, 338–345.
- Lindenmayer, D.B., Pierson, J.C., Barton, P. (Eds.), 2015. *Indicators and Surrogates of Biodiversity*. CSIRO Publishing, CRC Press, Melbourne, London.
- Mace, G.M., 2014. *Whose conservation?* *Science* 345, 1558–1560.
- McGeogh, M.A., 1998. The selection, testing and application of terrestrial insects as bioindicators. *Biol. Rev.* 73, 181–201.
- Miller, K.A., Miller, H.C., Moore, J.A., Mitchell, N.J., Cree, A., Allendorf, F.W., Sarre, S.D., Keall, S.N., Nelson, N.J., 2012. Securing the demographic and genetic future of tuatara through assisted colonization. *Conserv. Biol.* 26, 790–798.
- Moretti, M., Legg, C., 2009. Combining plant and animal traits to assess community functional responses to disturbance. *Ecography* 32, 299–309.
- Niemi, G.J., McDonald, M.E., 2004. Application of ecological indicators. *Annu. Rev. Ecol. Evol. Syst.* 35, 89–111.
- Noss, R.F., 1987. From plant communities to landscapes in conservation inventories: a look at the Nature Conservancy (USA). *Biol. Conserv.* 41, 11–37.
- Noss, R.F., 1990. Indicators for monitoring biodiversity – a hierarchical approach. *Conserv. Biol.* 4, 355–364.
- Pereira, H.M., Ferrier, S., Walters, M., Geller, G.N., Jongman, R.H.G., Scholes, R.J., Bruford, M.W., Brummitt, N., Butchart, S.H.M., Cardoso, A.C., Coops, N.C., Dulloo, E., Faith, D.P., Freyhof, J., Gregory, R.D., Heip, C., Höft, R., Hurtt, G., Jetz, W., Karp, D.S., McGeoch, M.A., Obura, D., Onoda, Y., Pettorelli, N., Reyers, B., Sayre, R., Scharlemann, J.P.W., Stuart, S.N., Turak, E., Walpole, M., Wegmann, M., 2013. *Essential biodiversity variables*. *Science* 339, 277–278.
- Rapport, D.J., Costanza, R., McMichael, A.J., 1998. Assessing ecosystem health. *Trends Ecol. Evol.* 13, 397–402.
- Rocha, F.C., Andrade, E.M., Lopes, F.B., 2015. Water quality index calculated from biological, physical and chemical attributes. *Environ. Monit. Assess.* 187, Art.4163.
- Schindler, D.E., Hilborn, R., Chasco, B., Boatright, C.P., Quinn, T.P., Rogers, L.A., Webster, M.S., 2010. Population diversity and the portfolio effect in an exploited species. *Nature* 465, 609–612.
- Schultz, P.W., 2011. Conservation means behavior. *Conserv. Biol.* 25, 1080–1083.
- Schwartz, M.K., Luikart, G., Waples, R.S., 2007. Genetic monitoring as a promising tool for conservation and management. *Trends Ecol. Evol.* 22, 25–33.
- Secretariat of the Convention on Biological Diversity, 2014. *Global Biodiversity Outlook 4*. Secretariat of the Convention on Biological Diversity, Montreal.
- Simberloff, D., 1998. Flagships, umbrellas, and keystones: is single-species management passe in the landscape era? *Biol. Conserv.* 83, 247–257.
- Stephens, P.A., Pettorelli, N., Barlow, J., Whittingham, M.J., Cadotte, M.W., 2015. Management by proxy? The use of indices in applied ecology. *J. Appl. Ecol.* 52, 1–6.
- Sutherland, W.J., Pullin, A.S., Dolman, P.M., Knight, T.M., 2004. The need for evidence-based conservation. *Trends Ecol. Evol.* 19, 305–308.
- Tallmon, D.A., Waples, R.S., Gregovich, D., Schwartz, M.K., 2012. Detecting population recovery using gametic disequilibrium-based effective population size estimates. *Conserv. Genet. Resour.* 4, 987–989.
- Triantis, K.A., Economo, E.P., Guilhaumon, F., Ricklefs, R.E., 2015. Diversity regulation at macro-scales: species richness on oceanic archipelagos. *Global Ecol. Biogeogr.* 24, 594–605.
- United Nations Convention to Combat Desertification (UNCCD) Advice on how best to measure progress on strategic objectives 1, 2 and 3 of The Strategy. UNCCD, Bonn, 2013. Available from <http://www.unccd.int/en/programmes/Science/Monitoring-Assessment/Documents/Decision22-COP11.pdf> (Accessed 16.12.15).
- US Fish and Wildlife Service (USFWS), Technical guidance on selecting species for landscape conservation, 2014. USFWS, Arlington VA, Available from <http://www.fws.gov/science/pdf/Draft-Technical-Guidance-on-Selecting-Species-for-Landscape-Scale-Conservation.pdf> (accessed 16.12.15).
- Walker, B., Salt, D., 2012. *Resilience Practice*. Island Press, Washington, DC.
- Westgate, M.J., Likens, G.E., Lindenmayer, D.B., 2013. Adaptive management of biological systems: a review. *Biol. Conserv.* 158, 128–139.
- Westgate, M.J., Barton, P.S., Lane, P.W., Lindenmayer, D.B., 2014. Global meta-analysis reveals low consistency of biodiversity congruence relationships. *Nat. Commun.* 5, 3899.
- Wikramanayake, E.D., Dinerstein, E., Robinson, J.G., Karanth, U., Rabinowitz, A., Olson, D., Mathew, T., Hedao, P., Conner, M., Hemley, G., Bolze, D., 2008. An ecology-based method for defining priorities for large mammal conservation: the tiger as case study. *Conserv. Biol.* 12, 865–878.